

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 471/04, A61K 31/47	A1	(11) International Publication Number: WO 98/55481 (43) International Publication Date: 10 December 1998 (10.12.98)
(21) International Application Number: PCT/EP98/03118 (22) International Filing Date: 27 May 1998 (27.05.98) (30) Priority Data: 197 23 161.6 3 June 1997 (03.06.97) DE 97108892.7 3 June 1997 (03.06.97) EP (34) Countries for which the regional or international application was filed: AT et al. (71) Applicant (for all designated States except US): BYK GULDEN LOMBERG CHEMISCHE FABRIK GMBH [DE/DE]; Byk-Gulden-Strasse 2, D-78467 Konstanz (DE). (72) Inventors (for all designated States except CA US): GUTTERER, Beate; Allensbacher Strasse 6b, D-78476 Allensbach (DE). ULRICH, Wolf-Rüdiger; Hebelstrasse 3, D-78464 Konstanz (DE). AMSCHLER, Hermann; Hohenhewenstrasse 19, D-78315 Radolfzell (DE). BEUME, Rolf; Bohlstrasse 13, D-78465 Konstanz (DE). HÄFNER, Dietrich; Beethovenstrasse 5, D-78464 Konstanz (DE). BÄR, Thomas; Blarerstrasse 16, D-78462 Konstanz (DE). MARTIN, Thomas; Sonnenbühlstrasse 73, D-78464 Konstanz (DE). KILIAN, Ulrich; Am Dachsberg 18, D-78479 Reichenau (DE). ELTZE, Manfred; Schützenstrasse 20, D-78462 Konstanz (DE). BRAND,		Ursula; Tägermoosstrasse 22, D-78462 Konstanz (DE). BOSS, Hildegard; Flurweg 3a, D-78464 Konstanz (DE). KLEY, Hans-Peter; Im Weinberg 3b, D-78476 Allensbach (DE). SCHUDT, Christian; Schützenstrasse 20, D-78462 Konstanz (DE). GOEBEL, Karl-Josef; Im Kirchental 12, D-78315 Radolfzell (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): FLOCKERZI, Dieter [DE/DE]; Ackerweg 26, D-78476 Allensbach (DE). HATZELMANN, Armin [DE/DE]; Alter Wall 3, D-78467 Konstanz (DE). (74) Common Representative: BYK GULDEN LOMBERG CHEMISCHE FABRIK GMBH; Byk-Gulden-Strasse 2, D-78467 Konstanz (DE). (81) Designated States: AL, AU, BA, BG, BR, CA, CN, CZ, EE, GE, HU, ID, IL, JP, KR, LT, LV, MK, MX, NO, NZ, PL, RO, SG, SI, SK, TR, UA, US, VN, YU, ZW, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: BENZONAPHTHYRIDINE (57) Abstract The invention concerns the compounds 8,9-Diethoxy-2-methyl-6-[4(p-toluenesulfonamido)phenyl]-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine and 9-Ethoxy-8-methoxy-2-methyl-6-[4-(p-toluenesulfonamido)phenyl]-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine and their therapeutic application.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Benzonaphthyridine

Field of application of the invention


The invention relates to novel benzonaphthyridines, a process for their preparation, their use and medicaments containing them. The compounds according to the invention are used in the pharmaceutical industry for the preparation of medicaments.

Known technical background

WO91/17991 describes under the title "New sulphonyl compounds" certain benzonaphthyridine derivatives, which are to be suitable for the treatment of airway disorders. In WO93/09780 and in DE-OS 4310050 the use of these benzonaphthyridine derivatives for the treatment of dermatoses, allergic rhinitis and conjunctivitis as well as of nasal polyps is described. For the compound (-)-cis-8,9-Dimethoxy-2-methyl-6-[4-(p-toluenesulfonamido)-phenyl]-1,2,3,4,4a,10b-hexahydrobenzo[c][1,6]naphthyridine, which is particularly emphasized in WO91/17991, WO93/09780 and DE 4310050, the WHO has proposed the INN Tolafentrine.

Description of the invention

The invention relates to compounds of the formula I



in which

R1 is ethoxy,

R2 is methoxy or ethoxy,
and the salts of these compounds.

The compounds of the formula I are chiral compounds having chiral centers in positions 4a and 10b. The invention therefore both comprises all conceivable pure diastereomers and pure enantiomers, and their mixtures in any mixing ratio, including the racemates. Preferred compounds of the formula I are those in which the hydrogen atoms in the positions 4a and 10b are cis to one another.

Particularly preferred are the compounds

(-)-cis-8,9-Diethoxy-2-methyl-6-[4-(p-toluenesulfonamido)-phenyl]-1,2,3,4,4a,10b-hexahydro-benzo-[c][1,6]naphthyridine and

(-)-cis-9-Ethoxy-8-methoxy-2-methyl-6-[4-(p-toluenesulfonamido)-phenyl]-1,2,3,4,4a,10b-hexahydro-benzo[c][1,6]naphthyridine

and the salts of these compounds.

Suitable salts for compounds of the formula I preferably are all acid addition salts. Particular mention may be made of the pharmacologically tolerable salts with the inorganic and organic acids customarily used in pharmacy. Examples of such suitable salts are water-soluble and water-insoluble acid addition salts with acids such as for example, hydrochloric acid, hydrobromic acid, phosphoric acid, nitric acid, sulfuric acid, acetic acid, citric acid, D-gluconic acid, benzoic acid, 2-(4-hydroxybenzoyl)benzoic acid, butyric acid, sulfosalicylic acid, maleic acid, lauric acid, malic acid, fumaric acid, succinic acid, oxalic acid, tartaric acid, embonic acid, stearic acid, toluenesulfonic acid, methanesulfonic acid or 3-hydroxy-2-naphthoic acid, where the acids are employed in salt preparation - depending on whether a mono- or polybasic acid is concerned and depending on which salt is desired - in an equimolar quantitative ratio or one differing therefrom.


Pharmacologically intolerable salts, which can be obtained, for example, as process products during the preparation of the compounds according to the invention on an industrial scale, are converted into pharmacologically tolerable salts by processes known to the person skilled in the art.

According to expert's knowledge the compounds of the invention as well as their salts may contain, e.g. when isolated in crystalline form, varying amounts of solvents. Included within the scope of the invention are therefore all solvates and in particular all hydrates of the compounds of formula I as well as all solvates and in particular all hydrates of the salts of the compounds of formula I.

The invention further relates to a process for the preparation of the compounds of the formula I, in which R1 and R2 have the meanings indicated above, and their salts.


The process is characterized in that

- a) compounds of the formula II



in which R₁ and R₂ have the abovementioned meanings, are reacted with a reactive derivative of p-toluenesulfonic acid, or in that

- b) compounds of the formula III



in which R₁ and R₂ have the abovementioned meanings, are subjected to a cyclocondensation reaction

and in that, if desired, compounds of the formula I obtained according to a) or b) are then converted into their salts, or in that, if desired, salts of the compounds of the formula I obtained according to a) or b) are then converted into the free compounds.

The reaction of compounds of the formula II with reactive derivatives of the p-toluenesulfonic acid (for example, a p-toluenesulfonic acid halide, particularly the acid chloride) is carried out in inert solvents in a manner known to the expert for the preparation of sulfonamides. The reaction is preferably carried out in the presence of an auxiliary base, such as for example, triethylamine or pyridine.

The cyclocondensation is carried out in a manner known per se to the person skilled in the art according to Bischler-Napieralski (e.g. as described in J. Chem. Soc., 1956, 4280-4282) in the presence of a suitable condensing agent, such as, for example, polyphosphoric acid, phosphorus pentachloride, phosphorus trichloride, phosphorus pentoxide, thionyl chloride or preferably phosphorus oxychloride, in a suitable inert solvent, e.g. in a chlorinated hydrocarbon such as chloroform, or in a cyclic hydrocarbon such as toluene or xylene, or another inert solvent such as acetonitrile, or without a further solvent using an excess of condensing agent, preferably at elevated temperature, in particular at the boiling temperature of the solvent or condensing agent used.

The described methods of preparation can be carried out analogously to the methods described in WO91/17991. The following examples serve to illustrate this.

Examples

1. (-)-cis-8,9-Diethoxy-2-methyl-6-[4-(p-toluenesulfonamido)-phenyl]-1,2,3,4,4a,10b-hexahydro-benzo[c][1,6]naphthyridine

A solution of 2.3 g p-toluenesulfonic acid chloride in 5 ml absolute dichloromethane is added dropwise to a solution of 3.5 g (-)-cis-6-(4-Aminophenyl)-8,9-diethoxy-2-methyl-1,2,3,4,4a,10b-hexahydrobenzo-[c][1,6]naphthyridine in 20 ml absolute pyridine, and the mixture is then stirred at room temperature for a further 3 h. After the evaporation of the solvents, the residue is extracted with dilute sodium hydroxide solution and dichloromethane. The organic phase is then washed with water, dried over sodium sulfate and concentrated. 5.4 g of the title compound are obtained as rough product, which is recrystallised twice in ethyl acetate/methanol. Yield: 4.3 g yellowish crystals. M.p. 267-268°C

EF: C₃₀H₃₅N₃O₄S, MW: 533.70

Optical rotation: $[\alpha]_D^{20} = -88.4^\circ$ (c=1, chloroform/methanol, 1+1)

$[\alpha]_{578\text{Hg}}^{20} = -93.2^\circ$ (c=1, chloroform/methanol, 1+1)

2. (-)-cis-9-Ethoxy-8-methoxy-2-methyl-6-[4-(p-toluenesulfonamido)-phenyl]-1,2,3,4,4a,10b-hexahydro-benzo[c][1,6]naphthyridine

2.52 g (-)-cis-3-(3-Ethoxy-4-methoxyphenyl)-1-methyl-4-[4-(p-toluenesulfonamido)-benzamido]-piperidine are heated to boiling under reflux for 5 h in 4.3 ml phosphorus oxychloride and 60 ml of acetonitrile. After distilling off the excess acetonitrile and phosphorus oxychloride, the residue is partitioned between dichloromethane and saturated sodium hydrogencarbonate solution. The organic phase is washed with water, dried over sodium sulfate and concentrated. After evaporation of the dichloromethane, the residue is purified over silica gel by chromatography. The main product fraction is separated and concentrated. The title compound is obtained after recrystallisation in ethyl acetate/diethyl ether (1:10) as faint yellow fine crystals. M.p. 207-219°C (unsharp, destruction and red colouring).

EF: C₂₉H₃₃N₃O₄S x 0.88 H₂O, MW: 535.49

Optical rotation: $[\alpha]_D^{20} = -65.1^\circ$ (c=1, methanol)

Starting compounds**A. (-)-cis-3-(3-Ethoxy-4-methoxyphenyl)-1-methyl-4-[4-(p-toluenesulfonamido)-benzamido]-piperidine**

The title compound is obtained by reaction of 1.36 g (-)-cis-4-Amino-3-(3-ethoxy-4-methoxyphenyl)-1-methylpiperidine with 4-(p-toluenesulfonamido)-benzoyl chloride [prepared from 1.5 g 4-(p-toluenesulfonamido)-benzoic acid and thionyl chloride] in dichloromethane under addition of triethylamine as auxiliary base. 2.65 g are obtained as solid foam. M.p. 100-105°C (the substance sticks together from about 93°C).

EF: C₂₉H₃₅N₃O₅S, MW: 537.68

Optical rotation: $[\alpha]_D^{20} = -69.6^\circ$ (c=1, methanol)

B. (-)-cis-4-Amino-3-(3-ethoxy-4-methoxyphenyl)-1-methylpiperidine dihydrochloride

The title compound is prepared analogously to the method described in DE 4217401, using rac-3-(3-Ethoxy-4-methoxyphenyl)-1-methylpiperid-4-one instead of rac-3-(3,4-Dimethoxyphenyl)-1-methylpiperid-4-one as starting material.

EF: C₁₅H₂₄N₂O₂ x 2HCl x 0.96 H₂O, MW: 354.52, [colourless crystals (isopropanol)], m.p. 252-254°C.

Optical rotation: $[\alpha]_D^{20} = -65.5^\circ$ (c=1, methanol)

C. (-)-cis-4-Amino-3-(3-ethoxy-4-methoxyphenyl)-1-methylpiperidine

The free base is prepared from the dihydrochloride (compound B) by treating with dilute sodium hydroxide solution and extraction with dichloromethane. It is used in the next reaction step without further purification.

Commercial utility

The compounds according to the invention have valuable pharmacological properties which make them commercially utilizable. As potent inhibitors of type 3, 4 and 5 of cyclic nucleotide phosphodiesterase (PDE3, PDE4 and PDE5), they are suitable on the one hand as bronchial therapeutics (for the treatment of airway obstructions on account of their dilating and cilium-stimulating action but also on account of their respiratory rate- and respiratory drive-increasing action), but on the other hand especially for the treatment of disorders of inflammatory nature, e.g. of the airways (asthma prophylaxis), of the skin, of the intestine, of the eyes and of the joints, which are mediated by mediators such as interferons, members of the tumor necrosis factor family, interleukins, chemokines, colony-stimulating factors, growth factors, lipid mediators (e.g., inter alia, PAF, platelet-activating factor), bacterial factors (e.g. LPS), immunoglobulins, oxygen free radicals and related free radicals (e.g. nitrogen monoxide NO), biogenic amines (e.g. histamine, serotonin), kinins (e.g. bradykinin), neurogenic mediators (such as substance P, neurokinin), proteins such as, for example, granular contents of leukocytes (inter alia cationic proteins of eosinophils) and adherent proteins (e.g. integrins). The compounds according to the invention have smooth muscle-relaxant action, e.g. in the region of the bronchial system, of the blood circulation, and of the efferent urinary passages. Furthermore they have a cilium-frequency increasing action, e.g. in the bronchial system.

In this context, the compounds according to the invention are distinguished by low toxicity, good human acceptance, great therapeutic breadth and the absence of significant side effects.

On account of their PDE-inhibiting properties, the compounds according to the invention can be employed as therapeutics in human and veterinary medicine, where they can be used, for example, for the treatment and prophylaxis of the following diseases: acute and chronic (in particular inflammatory and allergen-induced) airway disorders of various origin (bronchitis, allergic bronchitis, bronchial asthma); disorders with a reduction of the cilium activity or with increased demands on the ciliary clearance (bronchitis, mucoviscidose); dermatoses (especially of proliferative, inflammatory and allergic type) such as, for example, psoriasis (vulgaris), toxic and allergic contact eczema, atopic eczema, seborrheic eczema, lichen simplex, sunburn, pruritis in the anogenital area, alopecia areata, hypertrophic scars, discoid lupus erythematosus, follicular and widespread pyodermias, endogenous and exogenous acne, acne rosacea and other proliferative, inflammatory and allergic skin disorders; disorders which are based on excessive release of TNF and leukotrienes, i.e., for example, disorders of the arthritis type (rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis and other arthritic conditions), systemic lupus erythematosus, disorders of the immune system (AIDS), including AIDS-related encephalopathies, autoimmune disorders such as diabetes mellitus (Type I, autoimmune diabetes), multiple sclerosis and of the type virus-, bacteria- or parasite-induced demyelination diseases, cerebral malaria or Lyme's disease, shock symptoms [septic shock, endotoxin shock, Gram-negative sepsis, toxic shock syndrome and ARDS (adult respiratory distress syndrome)] and also generalized inflammations in the gastrointestinal region (Crohn's disease and ulcerative colitis); disorders which are

based on allergic and/or chronic, faulty immunological reactions in the region of the upper airways (pharynx, nose) and of the adjacent regions (paranasal sinuses, eyes), such as, for example, allergic rhinitis/sinusitis, chronic rhinitis/sinusitis, allergic conjunctivitis and also nasal polyps; and also disorders of the central nervous system such as memory disorders and Alzheimer's disease, candidiasis, leishmaniases and leprosy.

On account of their vasorelaxant activity, the compounds according to the invention can also be used for the treatment of high blood pressure disorders of various origin such as, for example, pulmonary high blood pressure and the concomitant symptoms associated therewith, for the treatment of erectile dysfunction or colics of the kidneys and the ureters in connection with kidney stones.

On account of their cAMP-increasing action, however, they can also be used for disorders of the heart which can be treated by PDE inhibitors, such as, for example, cardiac insufficiency, and also as anti-thrombotic, platelet aggregation-inhibiting substances.

The invention further relates to a method for the treatment of mammals including humans who are suffering from one of the abovementioned diseases. The method comprises administering a therapeutically effective and pharmacologically tolerable amount of one or more of the compounds according to the invention to the sick mammal.

The invention further relates to the compounds according to the invention for use in the treatment and/or prophylaxis of diseases, especially the diseases mentioned.

The invention also relates to the use of the compounds according to the invention for the production of medicaments which are employed for the treatment and/or prophylaxis of the diseases mentioned.

The invention furthermore relates to medicaments for the treatment and/or prophylaxis of the diseases mentioned and which contain one or more of the compounds according to the invention.

Advantageously, the substances according to the invention are also suitable for combination with other substances which bring about stimulation of cAMP, such as prostaglandins (PGE₂, PGI₂ and prostacyclin) and their derivatives, direct adenylate cyclase stimulators such as forskolin and related substances, or substances indirectly stimulating adenylate cyclase, such as catecholamines and adrenergic receptor agonists, in particular beta mimetics. In combination, on account of their cAMP degradation-inhibiting action, they in this case display a synergistic, superadditive activity. This comes to bear, for example, in their use in combination with PGE₂ for the treatment of pulmonary hypertension.

The medicaments are prepared by methods known per se familiar to the person skilled in the art. As medicaments, the compounds according to the invention (= active compounds) are either employed as such, or preferably in combination with suitable pharmaceutical auxiliaries, e.g. in the form of tab-

lets, coated tablets, capsules, suppositories, patches, emulsions, suspensions, gels or solutions, the active compound content advantageously being between 0.1 and 95%.

The person skilled in the art is familiar on the basis of his expert knowledge with the auxiliaries which are suitable for the desired pharmaceutical formulations. Beside solvents, gel-forming agents, ointments bases and other active compound excipients, it is possible to use, for example, antioxidants, dispersants, emulsifiers, preservatives, solubilizers or permeation promoters.

For the treatment of disorders of the respiratory tract, the compounds according to the invention are preferably also administered by inhalation. For this purpose, these are administered either directly as a powder (preferably in micronized form) or by atomization of solutions or suspensions which contain them. With respect to the preparations and administration forms, reference is made, for example, to the details in European Patent 163 965.

For the treatment of dermatoses, the compounds according to the invention are used in particular in the form of those medicaments which are suitable for topical application. For the production of the medicaments, the compounds according to the invention (= active compounds) are preferably mixed with suitable pharmaceutical auxiliaries and additionally processed to give suitable pharmaceutical formulations. Suitable pharmaceutical formulations which may be mentioned are, for example, powders, emulsions, suspensions, sprays, oils, ointments, fatty ointments, creams, pastes, gels or solutions.

The medicaments according to the invention are prepared by methods known per se. The dosage of the active compounds takes place in the order of magnitude customary for PDE inhibitors. Thus topical application forms (such as, for example, ointments) for the treatment of dermatoses contain the active compounds in a concentration of, for example, 0.1-99%. The dose for administration by inhalation is customarily between 0.1 and 3 mg per day. The customary dose in the case of systemic therapy (p.o. or i.v.) is between 0.01 and 10 mg/kg per day.

It is of particular interest for the present invention, that the compounds of formula I according to the invention clearly differ from the structurally closest compound of the state of the art - namely tolafentrine - in a surprising and for the person skilled in the art not foreseeable manner (Data are shown below in the chapter "Biological investigation").

The in-vitro data presented below for the compounds 1 and 2 (the numbers correspond to the numbers of the examples) show promise that the compounds 1 and 2 will have a clearly improved effectiveness in humans in comparison to tolafentrine. The higher potency with regard to the PDE4-inhibition indicates a considerably stronger antiinflammatory capacity, while the more distinct PDE3/PDE5-inhibition indicates a better broncholytic effectiveness.

Biological investigations

In the investigation of PDE4 inhibition at the cellular level, the activation of inflammatory cells has particular importance. An example which may be mentioned is the FMLP (N-formylmethionylleucyl-phenylalanine)-induced superoxide production of neutrophilic granulocytes, which can be measured as luminol-potentiated chemiluminescence [McPhail LC, Strum SL, Leone PA and Sozzani S, The neutrophil respiratory burst mechanism. In "Immunology Series" 1992, 57, 47-76; ed. Coffey RG (Marcel Decker, Inc., New York-Basel-Hong Kong)].

Substances which inhibit chemiluminescence, and/or cytokine secretion, and/or the secretion of inflammation-increasing mediators in inflammatory cells, like T-lymphocytes, monocytes, macrophages and granulocytes are those which inhibit PDE4 or PDE3 and PDE4. The latter isoenzyme of the phosphodiesterase families is particularly represented in granulocytes. Its inhibition leads to an increase in the intracellular cyclic AMP concentration and thus to the inhibition of cellular activation. PDE4 inhibition by the substances according to the invention is thus a central indicator of the suppression of inflammatory processes. (Giembycz MA, Could isoenzyme-selective phosphodiesterase inhibitors render bronchodilatory therapy redundant in the treatment of bronchial asthma? *Biochem Pharmacol* 1992, 43, 2041-2051; Torphy TJ et al., Phosphodiesterase inhibitors: new opportunities for treatment of asthma. *Thorax* 1991, 46, 512-523; Schudt C et al., Zardaverine: a cyclic AMP PDE3/4 inhibitor. In "New Drugs for Asthma Therapy", 379-402, Birkhäuser Verlag Basel 1991; Schudt C et al., Influence of selective phosphodiesterase inhibitors on human neutrophil functions and levels of cAMP and Ca; *Naunyn-Schmiedeberg's Arch Pharmacol* 1991, 344, 682-690; Tenor H and Schudt C, Analysis of PDE isoenzyme profiles in cells and tissues by pharmacological methods. In "Phosphodiesterase Inhibitors", 21-40, "The Handbook of Immunopharmacology", Academic Press, 1996; Hatzelmann A et al., Enzymatic and functional aspects of dual-selective PDE3/4-inhibitors. In "Phosphodiesterase Inhibitors", 147-160, "The Handbook of Immunopharmacology", Academic Press, 1996.

A. Methodology

1. Inhibition of the PDE isoenzymes

The PDE activity was determined according to Thompson et al. (1) with a few modifications (2). The test samples contained 40 mM tris HCl (pH 7.4), 5 mM $MgCl_2$, 0.5 μM cAMP or cGMP, [3H] cAMP or [3H]cGMP (about 50,000 cpm/sample), the PDE isoenzyme-specific additives described in greater detail below, the given concentrations of inhibitor and an aliquot of the enzyme solution in a total sample volume of 200 μl . Stock solutions of the compounds to be investigated in DMSO were prepared in concentrations such that the DMSO content in the test samples did not exceed 1% by volume - to avoid affecting the PDE activity. After preincubation at 37°C for 5 minutes, the reaction was started by addition of the substrate (cAMP or cGMP). The samples were incubated at 37°C for a further 15 min. The reaction was stopped by addition of 50 μl of 0.2N HCl. After cooling on ice for 10 minutes and addition of 25 μg of 5'-nucleotidase (snake venom from *Crotalus atrox*), incubation was carried out again for 10 min. at 37°C and the samples were then applied to QAE Sephadex A-25 columns. The columns were eluted with 2 ml of 30 mM ammonium formate (pH 6.0). The radioactivity of the eluate was measured and corrected by the corresponding blank values. The proportion of hydrolyzed nucleotide in no case exceeded 20% of the original substrate concentration.

PDE1 (Ca^{2+} /calmodulin-dependent) from bovine brain: the inhibition of this isoenzyme was investigated in the presence of Ca^{2+} (1 mM) and calmodulin (100 nM) using cGMP as a substrate (3).

PDE2 (cGMP-stimulated) from rats' hearts was purified chromatographically [Schudt et al. (4)] and investigated in the presence of cGMP (5 μM) using cAMP as a substrate.

PDE3 (cGMP-inhibited) and PDE5 (cGMP-specific) were investigated in homogenates of human blood platelets [Schudt et al. (4)] using cAMP or cGMP as a substrate.

PDE4 (cAMP-specific) was investigated in the cytosol of human polymorphonuclear leucocytes (PMNL) [isolated from leucocyte concentrates, see Schudt et al. (5)] using cAMP as a substrate. The PDE3 inhibitor motapizone (1 μM) was used in order to suppress the PDE3 activity emanating from contaminating blood platelets.

2. Inhibition of the formation of reactive oxygen species in human PMNL

The formation of reactive oxygen species determined by means of luminol-potentiated chemiluminescence (5) and the isolation of the PMNL from human blood (6) was carried out essentially as described in (5) and (6): equal-size portions (0.5 ml) of the cell suspension (10^7 cells/ml) were preincubated at 37°C for 5 min. in the absence or presence of the compounds to be investigated in a buffer solution containing 140 mM NaCl, 5 mM KCl, 10 mM HEPES, 1 mM $CaCl_2/MgCl_2$, 1 mg/ml of glucose, 0.05%

(w/v) BSA (bovine serum albumin), 10 μ M luminol and 4 μ M microperoxidase. Stock solutions of the compounds to be investigated in DMSO were prepared in such concentrations that the DMSO content - to avoid an effect on the PDE activity - in the test samples did not exceed 0.1% by volume. After preincubation, the test samples were additionally transferred to the measuring apparatus ["Multi-Biolumat" LB 9505C from Berthold (Wildbad, Germany)] before stimulation with the receptor agonist FMLP (N-formylmethionylleucylphenylalanine, 100 nM). The chemiluminescence was recorded continuously for 3 min.; the AUC values were calculated from this recording.

3. Statistics

The IC₅₀ values were determined from the concentration inhibition curves by nonlinear regression using the program GraphPad InPlot™ (GraphPad Software Inc., Philadelphia, USA).

4. References

- (1) Thompson W.J., Terasaki W.L., Epstein P.M. and Strada S.J., Assay of cyclic nucleotide phosphodiesterase and resolution of multiple molecular forms of the enzyme; Adv. Cycl. Nucl. Res. 1979, 10, 69-92
- (2) Bauer A.C. and Schwabe U., An improved assay of cyclic 3',5'-nucleotide phosphodiesterase with QAE Sephadex A-25; Naunyn-Schmiedeberg's Arch. Pharmacol. 1980, 311, 193-198
- (3) Gietzen K., Sadorf I. and Bader H., A model for the regulation of the calmodulin-dependent enzymes erythrocyte Ca²⁺-transport ATPase and brain phosphodiesterase by activators and inhibitors; Biochem. J. 1982, 207, 541-548.
- (4) Schudt C., Winder S., Müller B. and Ukena D., Zardaverine as a selective inhibitor of phosphodiesterase isoenzymes; Biochem. Pharmacol. 1991, 42, 153-162
- (5) Schudt C., Winder S., Forderkunnz S., Hatzelmann A. and Ullrich V., Influence of selective phosphodiesterase inhibitors on human neutrophil functions and levels of cAMP and Ca; Naunyn-Schmiedeberg's Arch. Pharmacol. 1991, 344, 682-690
- (6) Hatzelmann A. and Ullrich V., Regulation of 5-lipoxygenase activity by the glutathione status in human polymorphonuclear leukocytes; Eur. J. Biochem. 1987, 169, 175-184

B. Results

In Table 1 which follows, the inhibitory concentrations determined according to Section A1 [inhibitory concentrations as -log IC₅₀ (mol/l)] for the compounds according to the invention are indicated for various PDE isoenzymes. The numbers of the compounds correspond to the numbers of the examples.

Table 1

	PDE1	PDE2	PDE3	PDE4	PDE5
Tolafentrine	4.75	6.09	7.02	7.20	5.63
1	< 5	5.96	7.10	8.60	7.04
2	5.14	6.39	7.28	9.00	6.74

In table 2 below, the inhibitory concentrations determined according to Section A2 for tolafentrine and compound 1 are indicated for the FMLP-stimulated chemiluminescence in human PMNL.


Table 2

Inhibition of the FMLP-stimulated chemiluminescence in human PMNL in vitro by tolafentrine and compound 1 [inhibitory concentrations as $-\log IC_{50}$ (mol/l)].

Tolafentrine	6.07
1	7.39

Patent Claims

1. Compounds of the formula I



in which

R1 is ethoxy,

R2 is methoxy or ethoxy,

and the salts of these compounds.

2. A compound of the formula I with the chemical name (-)-cis-8,9-Diethoxy-2-methyl-6-[4-(p-toluenesulfonamido)-phenyl]-1,2,3,4,4a,10b-hexahydro-benzo[c][1,6]naphthyridine and the salts of this compound.
3. A compound of the formula I with the chemical name (-)-cis-9-Ethoxy-8-methoxy-2-methyl-6-[4-(p-toluenesulfonamido)-phenyl]-1,2,3,4,4a,10b-hexahydro-benzo[c][1,6]naphthyridine and the salts of this compound.
4. A medicament containing one or more compounds as claimed in claim 1 together with the customary pharmaceutical auxiliaries and/or excipients.
5. Compounds as claimed in claim 1 for use in the treatment of illnesses.
6. The use of compounds as claimed in claim 1 for the production of medicaments for the treatment of airway disorders.
7. The use of compounds as claimed in claim 1 for the production of medicaments for the treatment of high blood pressure disorders of various origins and the concomitant disorders connected therewith.

7. The use of compounds as claimed in claim 1 for the production of medicaments for the treatment of high blood pressure disorders of various origins and the concomitant disorders connected therewith.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/03118

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D471/04 A61K31/47

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 93 09780 A (BYK GULDEN LOMBERG CHEMISCHE FABRIK GMBH) 27 May 1993 cited in the application see the whole document ---	1,4
A	WO 91 17991 A (BYK GULDEN LOMBERG CHEMISCHE FABRIK GMBH) 28 November 1991 cited in the application see the whole document ---	1,4
A	EP 0 247 971 A (SANDOZ AG) 2 December 1987 see the whole document ---	1,4
A	DE 43 10 050 A (BYK GULDEN LOMBERG CHEMISCHE FABRIK GMBH) 14 October 1993 cited in the application see the whole document ---	1,4
-/--		



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

25 September 1998

Date of mailing of the international search report

02/10/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kyriakakou, G

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/03118

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 21 23 328 A (SANDOZ AG) 2 December 1971 see the whole document -----	1,4

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/03118

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9309780	A	27-05-1993	EP 0612247 A	31-08-1994
WO 9117991	A	28-11-1991	AU 651592 B	28-07-1994
			AU 7870191 A	10-12-1991
			CA 2082923 A	17-11-1991
			CN 1056691 A,B	04-12-1991
			CS 9101447 A	18-03-1992
			DE 4191109 A	01-04-1993
			EP 0528922 A	03-03-1993
			FI 925005 A	05-11-1992
			IL 98133 A	11-11-1994
			PT 97680 A	28-02-1992
			US 5346904 A	13-09-1994
EP 247971	A	02-12-1987	AU 599981 B	02-08-1990
			AU 7343787 A	03-12-1987
			DK 273187 A	30-11-1987
			FI 872360 A	30-11-1987
			JP 62294614 A	22-12-1987
			PT 84953 B	08-02-1990
DE 4310050	A	14-10-1993	NONE	
DE 2123328	A	02-12-1971	AT 326674 B	29-12-1975
			AT 412471 A	15-03-1975
			BE 767031 A	01-10-1971
			CA 970377 A	01-07-1975
			CH 527196 A	31-08-1972
			CS 162753 B	15-07-1975
			CS 167202 B	29-04-1976
			DK 134488 B	15-11-1976
			FI 52089 B	28-02-1977
			FR 2100649 A	24-03-1972
			GB 1361441 A	24-07-1974
			GB 1361442 A	24-07-1974
			NL 7106502 A	16-11-1971
			SE 387640 B	13-09-1976
			US 3899494 A	12-08-1975
			CH 527822 A	15-09-1972